



**Help protect
your child.**

At-a-glance guide to
childhood vaccines.

Why vaccines matter.

Thanks to widespread vaccination programs, several diseases that can infect our children have been eliminated. But there are still outbreaks of serious vaccine-preventable diseases that threaten the health of our children.

That's why immunizations against these 14 serious diseases are recommended by the Centers for Disease Control and Prevention (CDC). Each vaccine works differently to protect children from a wide range of diseases. So each one should be an essential part of your child's regularly scheduled wellness visits until your child is fully vaccinated.

Skipping vaccinations is a serious risk.

There are still outbreaks of serious, vaccine-preventable diseases. And that leaves babies and children who aren't vaccinated vulnerable to dangerous—even deadly—infections. If we stopped vaccinating, even the few cases we have in the United States could very quickly lead to tens or hundreds of thousands of cases.

Follow the CDC's full recommended vaccination schedule at the back of this brochure.



Help protect them when they're most vulnerable.

Making sure your child gets all the recommended vaccines is one of the most important steps you can take to help protect your child from infection. Receiving fewer than the recommended vaccinations leaves a child more vulnerable to catching serious diseases. And babies are more vulnerable because their immune systems are still developing. That's why most vaccinations are scheduled to start when babies are 2 months old.

Vaccinate on time. Every time.

Our children are at risk if parents skip vaccinations. So take the time to get familiar with 14 serious diseases and the vaccines that can help prevent them on the next few pages.

Answers to common questions parents ask.

Q. Do vaccines cause autism?

A. The National Institutes of Health funded a study that reviewed more than 20 major scientific studies and concluded that there is no evidence demonstrating vaccines cause autism.

Q. Are vaccines more dangerous than the diseases they protect against?

A. No. The safety and effectiveness of vaccines are tested in well-controlled scientific studies before they are approved by the US Food and Drug Administration.

Q. Are multiple vaccines too much for a child to receive in one day?

A. No. Babies' immune systems just get stronger as they grow and are exposed to more germs. And because the CDC recommends vaccination against 14 serious diseases, it's important to complete the full schedule to build your child's immunity.




Q. Do vaccines have side effects?

A. All vaccines can produce side effects. The most common include pain, redness, or tenderness where the injection is given. Though not common, serious side effects may occur, including allergic reactions; persistent, inconsolable crying; high fever; or seizures associated with fever. Be sure to talk to your doctor about any side effects.

Help protect your child.
Don't skip a doctor visit or a vaccine.

Turn the page for the CDC's full recommended vaccine schedule ▶

For Parents: Recommended immunizations for children from birth through 6 years old†



Is your family growing? To protect your new baby and yourself against whooping cough, get a Tdap vaccine. The recommended time is the 27th through 36th week of pregnancy. Talk to your doctor for more details.

Shaded boxes indicate the vaccine can be given during shown age range.

NOTE:

If your child misses a shot, you don't need to start over, just go back to your child's doctor for the next shot. Talk with your child's doctor if you have questions about vaccines.

FOOTNOTES:

- * Two doses given at least four weeks apart are recommended for children aged 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group.
- § Two doses of HepA vaccine are needed for lasting protection. The first dose of HepA vaccine should be given between 12 months and 23 months of age. The second dose should be given 6 to 18 months later. HepA vaccination may be given to any child 12 months and older to protect against HepA. Children and adolescents who did not receive the HepA vaccine and are at high-risk, should be vaccinated against HepA.

If your child has any medical conditions that put him at risk for infection or is traveling outside the United States, talk to your child's doctor about additional vaccines that he may need.

SEE BACK PAGE FOR MORE INFORMATION ON VACCINE-PREVENTABLE DISEASES AND THE VACCINES THAT PREVENT THEM.



For more information, call toll free **1-800-CDC-INFO** (1-800-232-4636) or visit www.cdc.gov/vaccines/parents



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention



American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

CDC-recommended immunization schedule, 2018

† This is a resource intended for parents or caregivers; there is a more detailed healthcare professional version of the schedule available on the CDC website.

Vaccinate on time. Every time.

Don't leave your child vulnerable to these vaccine-preventable diseases.

Chickenpox

Vaccine: Varicella



Chickenpox is a virus that can cause an itchy, blistery rash all over the body, and is generally accompanied by a fever and drowsiness. It is transmitted from person to person through the air or by contact with fluid from the rash. Serious complications may include skin infections, pneumonia, encephalitis (inflammation of the brain), and even death.

Diphtheria

Vaccine: DTaP



Diphtheria is caused by bacteria that live in an infected person's mouth or throat, and can cause a sore throat or fever, and may obstruct breathing. Sneezing or coughing can spread these bacteria from person to person. Serious complications may include heart failure, paralysis, and death.

Hib

Vaccine: Hib



Hib (*Haemophilus influenzae type B*) is caused by bacteria that are spread through the air by coughing or sneezing. It can cause ear infections and serious throat swelling. If Hib bacteria enter a person's bloodstream, they can cause meningitis, pneumonia, and other problems. Serious complications may include permanent brain damage and death.



Hepatitis A

Vaccine: HepA

Hepatitis A is a virus found mostly in bowel movements, and is spread by personal contact or through contaminated food or water. It can cause liver disease, which can result in stomach pain, vomiting, fever, and other problems. Serious complications may include liver failure that leads to death.



Hepatitis B

Vaccine: HepB

Hepatitis B is a virus that can cause liver disease and yellow skin or eyes (jaundice). It can spread through contact with infected blood or other body fluids, or from mother to baby at birth. Serious complications may include chronic liver disease, cirrhosis (scarring of the liver), liver cancer, and death.



Flu

Vaccine: Flu

Flu (Influenza) is a virus that is spread from person to person by droplets from coughing, sneezing, or talking, or from surfaces that have the virus on them. The flu can cause fever, sore throat, cough, chills, and muscle aches. Serious complications may include pneumonia, inflammation of the heart, and death.



Measles

Vaccine: MMR

Measles is a virus that can cause a rash all over the body, fever, runny nose, and cough. It is very contagious and is spread from person to person through coughing, sneezing, and even breathing. Serious complications may include pneumonia, seizures, permanent brain damage, and even death.



Mumps

Vaccine: MMR

Mumps is a virus that can cause fever, headache, and inflammation of the salivary glands, which leads to swelling of the cheeks and jaws. Person-to-person transmission occurs through the air. Serious complications may include meningitis, and occasionally encephalitis (inflammation of the brain) or deafness, and even death.



Pertussis

Vaccine: DTaP

Pertussis (Whooping cough) is caused by bacteria that are spread from person to person through the air. The disease can cause violent coughing spells that can affect eating, drinking, and even breathing. Serious complications may include pneumonia, seizures, encephalopathy (brain infection), and death.



Polio

Vaccine: IPV

Polio is a very contagious virus that can cause paralysis. Most infected people show no symptoms. It is spread through contact with the stool of an infected person or by droplets from a sneeze or cough. Serious complications may include weakness in arms or legs (or both), paralysis, and death.



Pneumococcal disease

Vaccine: PCV

Pneumococcal disease is caused by bacteria that are spread by airborne droplets, or by direct contact with infected saliva or mucus. Invasive disease can cause cough, fever and chills, chest pain, ear infections, blood infections, and difficulty breathing. Serious complications may include bacterial meningitis, which may lead to death.



Rotavirus

Vaccine: RV

Rotavirus spreads easily by hands, diapers, or objects that have a small amount of infected stool on them. This virus causes severe diarrhea, vomiting, and fever. Serious complications may include severe diarrhea leading to extreme dehydration, which can cause death.



Rubella

Vaccine: MMR

Rubella (German measles) is an airborne virus that causes swollen glands, a slight fever, rash, and occasionally arthritis-like symptoms. It is a mild disease in children. Serious complications are found in babies if the mother has been infected during pregnancy. The baby may be born deaf or blind, with a damaged heart or small brain, or be mentally impaired.



Tetanus

Vaccine: DTaP

Tetanus (Lockjaw) is a bacterial disease that enters through the skin from deep cuts and puncture wounds. Tetanus may cause headache, irritability, and spasms in the jaw muscles. Serious complications may include inability to swallow, muscle cramps so strong that they can break a child's bones, and often death.

Be sure to follow the full vaccination schedule recommended by the CDC.

More resources for you. Log on today.

- **Customizable online vaccination schedule**
www.vaccinecalendar.com

The organizations listed below have created helpful websites that provide additional information about childhood vaccines.

- **Centers for Disease Control and Prevention***
www.cdc.gov/vaccines/parents

- **American Academy of Pediatrics***
www.healthychildren.org

- **American Academy of Family Physicians***
www.familydoctor.org

*These websites are neither owned nor controlled by Pfizer. Pfizer is not responsible for the content or services of these sites.

